

EMERGENCY KIT

This emergency pack should contain one of each of the following:-

- ◆ Copy of the Emergency letter.
- ◆ Adrenal Insufficiency Card.
- ◆ Alcohol Wipe and cotton wool ball.
- ◆ Syringe & Needle.
- ◆ Ampoule of hydrocortisone for injection Efcortisol® recently renamed as (Hydrocortisone Sodium Phosphate) (100mg in 1ml) or Solu-Cortef® 100mgs in 2 mls. Solu-Cortef® comes in powder form with 1 vial of sterile water (2mls to mix), see insert on how to mix.
- ◆ How to give an injection of Efcortisol® (Hydrocortisone Sodium Phosphate) information booklet.
- ◆ Tube of Glucogel® (used to be known as HypoStop) For full instructions on how to use Glucogel® see Emergency Travel Leaflet.
- ◆ Oral Solution of Dioralyte in case of diarrhoea.

Remember that:-

- If you/your child is ill we now recommend ***an extra double/triple*** dose at ***04:00 am***. *This dose should be the same as the morning double/triple dose* and give the morning double/triple dose at 07:00 am.
- Glucogel® should only be used when you/your child is showing signs of low blood sugar levels.
- Glucogel® should be used **AFTER** you/your child has had the hydrocortisone injection.
- It is advisable that you take several ampoules of Efcortisol (Hydrocortisone Sodium Phosphate) with you in case an ampoule gets broken, or crushes as you try to break it.

DEALING WITH DIARRHOEA

- Diarrhoea is a particular problem due to the fluid losses.
- Oral rehydration solutions such as Dioralyte should be used.
- Make sure you/your child is passing urine regularly.
- Seek medical advice early especially:-
 - ~ If there is a fever.
 - ~ Blood in the diarrhoea.
 - ~ Your child becomes confused.
 - ~ The diarrhoea does not stop after 24 hours.
- Do not use anti diarrhoeal drugs.

REMEMBER

You will need a letter of authorisation to allow you to carry the injection kit with you through security checks and customs. Please remember to allow your endocrinologist enough time to organise the letter before you travel.

**THE EMERGENCY LETTER
GENERAL ANAESTHETIC PROTOCOL ARE
ACCESSIBLE FROM OUR WEBSITE**

<http://www.cahisus.co.uk/>

USEFUL CONTACT NUMBER

Hospital for Tropical Diseases Travel Clinic
Mortimer Market (off Tottenham Court Road)

London WC1E 6AU


Tel: 020 7388 9600

Healthline (charges 50p/minute): 09061 33 77 33

This leaflet is a joint production between
Professor Peter Hindmarsh and Kathy Geertsma

The series editor is
Professor Peter Hindmarsh
Professor of Paediatric Endocrinology
and Divisional Clinical Director for Paediatrics
and Adolescents at UCLH
<http://www.cahisus.co.uk>

"The ideas expressed are independent of the editors' affiliations. Data provided is from current literature and should always be discussed with your endocrinologist"



Hypopituitarism

TRAVEL ADVICE

TIME ZONES

Professor Peter Hindmarsh

Air travel means that patients receiving hydrocortisone treatment will need to face changes to their dosing schedule. This information leaflet tells you how to adjust doses for different travel destinations. Always carry your medication with you in your hand luggage.

AIR TRAVEL AND TIME ZONES

Increasing air travel means that many patients receiving hydrocortisone treatment have to face changes to their dosing schedule. This information sheet tells you how to adjust doses for different travel destinations.

You might also need advice on additional vaccinations or tablets, so a call to the Travel Clinic or as most GP's run a travel clinic, an appointment to discuss what is required is a good idea.

PLEASE REMEMBER

- You/your child should wear a Medic-Alert bracelet or equivalent at all times.
- Take a copy of the Emergency letter with you, remember it can also be accessed from our website
- Take you/your child's Adrenal Insufficiency Card with you and make sure it is up to date.
- Carry your/your child's supply of hydrocortisone for emergency use in your hand luggage.
- Make sure all medication is in date and allow plenty of time to get a new supply if it has expired.
- Ensure you have enough supplies of everything for the length of your holiday plus a few days more.
- Also make sure you have an extra supply in case of illness.
- Getting medicines abroad can sometimes be difficult and the formulations are not always the same.

EMERGENCY PACK

- Keep the entire emergency pack together, in a safe place where children cannot reach it.
- Keep it at room temperature, out of direct sunlight or heat.
- The injection ampoules do not need to be kept in the fridge.

TIME ZONES

For travel within Europe this is not a problem but longer journey times may need some adjustment to the timings of doses. See chart of countries and dose adjustments.

THYROXINE AND GROWTH HORMONE

If you are taking thyroxine tablets or Growth Hormone injections then they should continue as once per day as usual. Simply adopt the local time that you would normally take the medication at home.

MEDICATION CHANGES DURING TRAVEL TWICE DAILY HYDROCORTISONE THERAPY

Africa (Including South Africa)

Journey to and from Africa

Note:- Within a 1 to 3 hour time difference, no change is required.

Europe

Journey to and from Europe

Note:-
Within a 1 to 3 hour time shift, no change is required

Far East, Australia or New Zealand

Journey to and from Far East, Australia or New Zealand

Note:-
For these journeys, it is better to switch to an 8-hourly treatment programme
Normal daily schedule up to departure
Switch to 8-hourly schedule for flight duration
Continue 8-hourly schedule until morning after arrival
Go back to normal dose schedule

Middle East or India

Journey to Middle East or India

Evening dose as usual
Half morning dose on arrival
Normal dose schedule from the morning of arrival day

Journey from Middle East or India

Evening dose as usual
Repeat evening dose when boarding plane
Usual morning dose on arrival in UK
Normal evening dose
Normal dose schedule the following day

United States of America

Journey to USA

Morning dose as usual
Half morning dose on arrival
Evening dose before going to bed
Normal dose schedule the following day

Journey from USA

Evening dose as usual
Normal dose schedule the following day

MEDICATION CHANGES DURING TRAVEL THREE TIME DAILY HYDROCORTISONE THERAPY

Africa (Including South Africa)

Journey to and from Africa

Note:- Within a 1 to 3 hour time difference, no change is required.

Europe

Journey to and from Europe

Note:-
Within a 1 to 3 hour time shift, no change is required

Far East, Australia or New Zealand

Journey to and from Far East, Australia or New Zealand

Note:-
No significant changes as dose schedule is already 8-hourly

Middle East or India

Journey to Middle East or India

Evening dose as usual
Half morning dose on arrival
Normal dose schedule from the morning of arrival day

Journey from Middle East or India

Repeat evening dose on boarding plane
Usual morning dose on arrival in UK
Second dose late afternoon
Normal evening dose
Normal dose schedule the following day

United States of America

Journey to USA

Morning dose as usual
Half morning dose on arrival
Evening dose before going to bed
Normal dose schedule the following day

Journey from USA

Evening dose as usual
Normal dose schedule the following day